

PHYTOCHEMICALS AND BIOLOGICAL ACTIVITIES OF *ARTOCARPUS*
FULVICORTEX JARRETT

NUR AZLIN BINTI OTHMAN

UNIVERSITI TEKNOLOGI MALAYSIA

PHYTOCHEMICALS AND BIOLOGICAL ACTIVITIES OF *ARTOCARPUS*
FULVICORTEX JARRETT

NUR AZLIN BINTI OTHMAN

A thesis submitted in fulfillment of the
requirements for the award of the degree of
Master of Science (Chemistry)

Faculty of Science
Universiti Teknologi Malaysia

2010

Dedicated to..

My parents, Othman bin Taib and Rahimah bte Mohamed

My husband, Mohd Nazrul bin Mohd Amin

My sisters and brothers

ACKNOWLEDGEMENTS

Foremost, I would like to express my sincere gratitude to my supervisor, Dr. Shajarahtunnur bte Jamil for the continuous support of my study and research, for her patience, motivation, enthusiasm, and immense knowledge. Her guidance helped me in all the time of research and writing of this thesis.

I am deeply grateful to my co-supervisor, Dr. Muhamed Taher and his student, Mr. Osama from Kuliyyah of Pharmacy, International Islamic University Malaysia for their important help throughout the cytotoxicity experimental work.

My sincere thanks also goes to Ministry of Science, Technology, and Innovation (MOSTI) for the financial support *via* vote number 79132.

During this work, I have collaborated with many people in the Department of Chemistry for whom I have great regard, and I wish to extend my warmest thanks to all those who have helped me with my work.

I also extend my thanks to Natural Product Research Group: Dr. Emrizal, Fazlin, Hafiz, Fariz, Norakmalazura, Syamil, Yau, Ummu, Salihin, Safina, Noraini, Saidah, and Raihan for the friendship and invaluable assistance.

Last but not least, I owe the most special thanks to my parents, husband and siblings for their unflagging love, support, and pray for my success.

PREFACE

This thesis is the result of my work carried out in the Department of Chemistry, Universiti Teknologi Malaysia and Kuliyyah of Pharmacy, International Islamic University Malaysia between December 2007 and December 2009 under the supervision of Dr. Shajarahtunnur bte Jamil and Dr. Muhamed Taher. Parts of my work described in this thesis have been reported in the following publications:

1. Nur Azlin Othman, Shajarahtunnur Jamil, and Muhamed Taher. Phytochemicals with Cytotoxic Activity from *Artocarpus fulvicortex*. Paper presented at the International Conference on Molecular Chemistry 2008 at Universiti Malaya, Kuala Lumpur. November 25-27, 2008.
2. Nur Azlin Othman, Shajarahtunnur Jamil, and Muhamed Taher. Antioxidant Activity of Flavonoids Isolated from the Leaves of *Artocarpus fulvicortex*. Poster presented at the International Conference and Workshops on Basic and Applied Science & Regional Annual Fundamental Science Seminar 2009 at The Zone Regency Hotel, Johor Bahru. June 3-4, 2009.
3. Nur Azlin Othman, Shajarahtunnur Jamil, and Muhamed Taher. Cytotoxic Activity of Flavonoids Isolated from the Barks of *Artocarpus fulvicortex*. Paper presented at the Junior Chemist Colloquium 2009 at Universiti Malaysia Sarawak, Kota Samarahan. July 1-2, 2009.

ABSTRACT

Phytochemical studies of the leaves and barks of *Artocarpus fulvicortex* Jarrett have been carried out. Cold extraction method was applied to obtain the crude extracts. Twelve compounds, comprising of chalcones, flavones, and triterpenes were successfully isolated from this plant. The structures of these pure compounds were determined on the basis of spectral studies including nuclear magnetic resonance, infrared, ultraviolet spectroscopies, mass spectrometry as well as by comparison with literature data. Multiple purification of the leaves crude extracts yielded a new flavonoid, identified as 5-hydroxy-2",2"-dimethylpyrano-[5",6",11",12":6,7,3',4'] flavone together with known compounds i.e. carpachromene, cycloartocarpesin, 2'-hydroxy-4,4',6'-trimethoxychalcone, norartocarpetin, friedelin, lupeol, and β -sitosterol. Five known compounds, characterized as artobiloxanthone, artonin E, catechin, lupeol 3-acetate, and friedelin were isolated from the barks crude extracts. Evaluations on biological activities were performed by using antibacterial, antioxidant, and cytotoxic assays on the pure compounds. Artobiloxanthone and artonin E were found to have antibacterial activity against *Pseudomonas aeruginosa* and *Bacillus subtilis* with MIC values in the range of 62.5-250 $\mu\text{g/mL}$, while MBC values were at 125-500 $\mu\text{g/mL}$. Catechin and artonin E showed high ability to act as free radical scavenger with SC_{50} values of 19.4 $\mu\text{g/mL}$ and 38.2 $\mu\text{g/mL}$ respectively. The cytotoxic activity was done by using 3-(4,5-dimethylthiazol-2-yl)-2-5-diphenyltetrazolium bromide (MTT) assay on esophagus cancer (TE-1), lung cancer (A549), breast cancer (MCF-7), cervix cancer (C33A), and colon cancer (CaCO₂) cell lines. Artonin E displayed significant effect against esophagus cancer (TE-1) and colon cancer (CaCO₂) cell lines with the percentage of cell survival of 24.72% and 47.29% at 11 $\mu\text{g/mL}$. Carpachromene, cycloartocarpesin, 2'-hydroxy-4,4',6'-trimethoxychalcone, friedelin, and lupeol showed moderate activity against colon cancer (CaCO₂) cell line with percentage of cell survival of 52 to 70%.

ABSTRAK

Kajian fitokimia ke atas daun dan kulit batang *Artocarpus fulvicortex* Jarrett telah dilakukan. Pengekstrakan secara sejuk telah dijalankan bagi mendapatkan ekstrak mentah. Dua belas sebatian yang terdiri daripada kalkon, flavon, dan triterpena telah berjaya diasingkan daripada tumbuhan ini. Struktur bagi sebatian tulen dikenalpasti berdasarkan kajian spektroskopi yang terdiri daripada spektroskopi resonans magnetik nukleus, infra merah, ultra lembayung, spektrometri jisim, dan juga perbandingan dengan data dari kajian terdahulu. Penulenan yang berulang kali terhadap ekstrak mentah daun telah memberikan flavonoid baru, dikenalpasti sebagai 5-hidroksi-2'',2''-dimetilpirano-[5'',6'',11'',12'':6,7,3,4']flavon bersama dengan sebatian yang diketahui, iaitu karpakromen, sikloartokarpesin, 2'-hidroksi-4,4',6'-trimetoksikalkon, norartokarpetin, friedelin, lupeol, dan β -sitosterol. Lima sebatian yang diketahui dan dikenalpasti sebagai artobiloxanton, artonin E, katekin, lupeol 3-asetat, dan friedelin telah diasingkan daripada ekstrak mentah kulit batang. Kajian bioaktiviti telah dilakukan ke atas sebatian tulen dengan menggunakan cerakanan antibakteria, antioksidan, dan sitotoksik. Artobiloxanton dan artonin E didapati mempunyai aktiviti antibakteria terhadap *Pseudomonas aeruginosa* and *Bacillus subtilis* dengan nilai MIC pada anggaran 62.5-250 $\mu\text{g/mL}$, sementara nilai MBC pada 125-500 $\mu\text{g/mL}$. Katekin dan artonin E menunjukkan keupayaan yang tinggi sebagai perencat radikal bebas dengan nilai 19.4 $\mu\text{g/mL}$ and 38.2 $\mu\text{g/mL}$. Kajian sitotoksik telah dilakukan dengan menggunakan cerakanan 3-(4,5-dimetiltiazol-2-il)-2-5-difeniltetrazolium bromid (MTT) ke atas sel kanser esofagus (TE-1), kanser paru-paru (A549), kanser payudara (MCF-7), kanser serviks (C33A), dan kanser kolon (CaCO₂). Artonin E memperlihatkan kesan yang signifikan terhadap sel kanser esofagus (TE-1) dan kanser kolon (CaCO₂) dengan peratusan sel kanser hidup 24.72% dan 47.29% pada 11 $\mu\text{g/mL}$. Karpakromen, sikloartokarpesin, 2'-hidroksi-4,4',6'-trimetoksikalkon, friedelin, dan lupeol menunjukkan aktiviti sederhana terhadap sel kanser kolon (CaCO₂) dengan peratusan sel kanser hidup antara 52 hingga 70%.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	TITLE PAGE	i
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENTS	iv
	PREFACE	v
	ABSTRACT	vi
	ABSTRAK	vii
	TABLE OF CONTENTS	viii
	LIST OF TABLES	xii
	LIST OF SCHEMES	xiv
	LIST OF FIGURES	xv
	LIST OF ABBREVIATIONS	xvi
	LIST OF APPENDICES	xix
1	INTRODUCTION	
1.1	General Introduction	1
1.2	Family Moraceae	2
1.3	Genus <i>Artocarpus</i>	3
	1.3.1 <i>Artocarpus fulvicortex</i> Jarrett	4
1.4	Problem Statement	4
1.5	Research Objectives	4
1.6	Scope of Study	5

2 LITERATURE REVIEW

2.1	Review on Phytochemicals and Biological Activities of <i>Artocarpus</i> Species	6
2.1.1	Isoprenylflavonoids and Derivatives	7
2.1.2	Stilbenes and 2-Arylbenzofuran Derivatives	18
2.1.3	Geranylated Phenolic Constituents	21
2.1.4	Natural Diels-Alder Type Adducts	25
2.1.5	Terpenoids	
2.2	Biosynthesis Pathway of Flavonoids	28

3 RESULTS AND DISCUSSION

3.1	Phytochemicals of the Leaves of <i>Artocarpus fulvicortex</i> Jarrett	30
3.1.1	Carpachromene (107)	31
3.1.2	Cycloartocarpesin (108)	34
3.1.3	5-Hydroxy-2'',2''-dimethylpyrano-[5'',6'',11'',12'':6,7,3',4']flavone (106)	36
3.1.4	2'-Hydroxy-4,4',6'-trimethoxychalcone (109)	39
3.1.5	Norartocarpetin (48)	42
3.1.6	Friedelin (110)	44
3.1.7	Lupeol (111)	46
3.1.8	β -sitosterol (100)	47
3.2	Phytochemicals of the Barks of <i>Artocarpus fulvicortex</i> Jarrett	49
3.2.1	Artonin E (22)	49
3.2.2	Artobiloxanthone (21)	52
3.2.3	Catechin (112)	54
3.2.4	Lupeol 3-acetate (101)	57
3.2.5	Friedelin (110)	58
3.3	Bioactivity Studies of <i>Artocarpus fulvicortex</i> Jarrett	58
3.3.1	Introduction and the Principle of Antibacterial Activity	58
	3.3.1.1 Antibacterial Activity of <i>A. fulvicortex</i> Jarrett	59
3.3.2	Introduction and the Principle of Antioxidant Activity	62
	3.3.2.1 Free Radical Scavenging (DPPH)	63

	3.3.2.2 Antioxidant Activity of <i>A. fulvicortex</i> Jarrett	63
	3.3.3 Introduction and the Principle of Cytotoxic Activity	65
	3.3.3.1 Cytotoxic Activity of <i>A. fulvicortex</i> Jarrett	65
4	EXPERIMENTAL	
4.1	General Experimental Procedures	67
4.2	Chemicals	68
4.3	Plant Materials	68
4.4	Isolation of Phytochemicals from the Leaves of <i>A. fulvicortex</i> Jarrett	68
4.4.1	Friedelin (110)	69
4.4.2	5-hydroxy-2",2"-dimethylpyrano-[5",6",11",12": 6,7,3',4']flavone (106)	70
4.4.3	Lupeol (111)	70
4.4.4	2'-Hydroxy-4,4',6'-trimethoxychalcone (109)	71
4.4.5	β -Sitosterol (100)	72
4.4.6	Carpachromene (107)	72
4.4.7	Cycloartocarpesin (108)	73
4.4.8	Norartocarpetin (48)	74
4.5	Isolation of Phytochemicals from the Barks of <i>A. fulvicortex</i> Jarrett	74
4.5.1	Lupeol 3-acetate (101)	75
4.5.2	Artonin E (22)	76
4.5.3	Artobiloxanthone (21)	76
4.5.4	Catechin (112)	77
4.5.5	Friedelin (110)	78
4.6	Bioactivity Studies	78
4.6.1	Instrumentations for Bioactivity Assay	78
4.6.2	Chemicals for Bioactivity Assay	79
4.6.3	Microorganisms and Cell Lines for Bioactivity Assay	79
4.6.4	Antibacterial Assay	80
	4.6.4.1 Microorganisms and Culture Media	80
	4.6.4.2 Disc Diffusion Method	80
	4.6.4.3 Minimum Inhibition Concentration (MIC)	81

4.6.4.4 Minimum Bactericidal Concentration (MBC)	81
4.6.5 Antioxidant Assay [Free Radical Scavenging Activity (DPPH)]	83
4.6.6 Cytotoxic Assay	84
5 CONCLUSIONS AND RECOMMENDATIONS	
5.1 Phytochemical Study	86
5.2 Biological Studies	86
5.3 Recommendations	87
REFERENCES	88
APPENDICES	98

LIST OF TABLES

TABLE NO.	TITLE	PAGE
1.1	<i>Artocarpus</i> Species Found in Malaysia	3
3.1	¹ H NMR, ¹³ C NMR, and COSY Data of Carpachromene (107)	32
3.2	NMR Data of Cycloartocarpesin (108)	36
3.3	¹ H NMR, ¹³ C NMR, and COSY Data of 5-hydroxy-2'',2''-dimethylpyrano-[5'',6'',11'',12'':6,7,3',4']flavone (106)	38
3.4	NMR Data of 2'-Hydroxy-4,4',6'-trimethoxychalcone (109)	41
3.5	¹ H NMR, ¹³ C NMR, and COSY Data of Norartocarpetin (48)	44
3.6	¹ H NMR, ¹³ C NMR, and COSY data of Artonin E (22)	50
3.7	NMR Data of Artobiloxanthone (21)	54
3.8	¹ H NMR, ¹³ C NMR, and HMBC Data of Catechin (112)	56
3.9	Antibacterial Activity of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett	60
3.10	Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett	61
3.11	Antioxidant Activity of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett	64
3.12	Cytotoxic Activity of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett by MTT Assay	66
4.1	Results of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett	82
4.2	Results of Antioxidant Activity of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett Measured by UV Spectrophotometric Method	84

4.3	Cytotoxic Activity of the Isolated Compounds of <i>A. fulvicortex</i> Jarrett by MTT Assay	85
-----	---	----

LIST OF SCHEMES

SCHEME NO.	TITLE	PAGE
3.1	The EIMS Fragmentation Pattern of Compound (107)	33
3.2	The Loss of Methyl Group from Compound (106)	37
3.3	The EIMS Fragmentation Pattern of Compound (109)	42
3.4	The EIMS Fragmentation Pattern of Compound (22)	51
3.5	The Loss of Methyl Group from Compound (21)	52
3.6	The EIMS Fragmentation Pattern of Compound (112)	56

LIST OF FIGURES

FIGURES NO.	TITLE	PAGE
2.1	Biosynthesis Pathway of Flavonoids	29
3.1	The Reduction Reaction of DPPH	63
4.1	The Arrangement of the Sample Discs and Control Discs in Petri Dish	81

LIST OF ABBREVIATIONS

Abs	Absorbance
AlCl ₃	Aluminium trichloride
br	broad
CC	Column Chromatography
COSY	Correlation Spectroscopy
¹³ C	Carbon-13
CDCl ₃	Deuterated chloroform
CD ₃ COCD ₃	Deuterated acetone
DPPH	2,2-Diphenyl-1-picrylhydrazyl
d	doublet
dd	doublet of doublet
ddd	doublet of doublet of doublet
dt	doublet of triplet
DEPT	Distortionless Enhancement by Polarization Transfer
DMSO	Dimethylsulfoxide
EtOAc	Ethyl acetate
EIMS	Electron Impact Mass Spectrometry
Et ₂ O	Diethyl ether
EtOH	Ethanol
FABMS	Fast Atom Bombardment Mass Spectrometry
GC	Gas Chromatography
GC-MS	Gas Chromatography-Mass Spectrometry
¹ H	Proton
HMBC	Heteronuclear Multiple Bond Correlation
HMQC	Heteronuclear Multiple Quantum Coherence
Hz	Hertz

HCl	Hydrochloric acid
IR	Infrared
IC ₅₀	Inhibition Concentration at 50%
<i>J</i>	coupling constant
KBr	Potassium bromide
KI	Kovats Index
lit.	Literature
<i>LWT</i>	Lebensm.-Wiss. u.-Technol / Food Science and Technology
MIC	Minimum Inhibition Concentration
MBC	Minimum Bactericidal Concentration
MS	Mass Spectrometry
mM	milimolar
<i>m/z</i>	mass to charge ion
MeOH	Methanol
mp	melting point
MgSO ₄	Magnesium sulphate
MHz	Megahertz
m	multiplet
NMR	Nuclear Magnetic Resonance
nm	nanometer
NaOH	Sodium hydroxide
NaCl	Sodium chloride
Ph	Phenyl
PE	Petroleum ether
ppm	parts per million
q	quartet
RDA	Retro Diels-Alder rearrangement
R _f	retention factor
SD	Standard Deviation
SFE	Supercritical Fluid Extraction
s	singlet
sh	shoulder
t	triplet
tr	trace

TLC	Thin Layer Chromatography
UV	Ultraviolet
VLC	Vacuum Liquid Chromatography
μM	micromolar
δ	chemical shift
$[\alpha]_{\text{D}}$	specific rotation
c	concentration

LIST OF APPENDICES

APPENDIX	TITLE	PAGE
1	Infrared spectrum of carpachromene (107)	98
2	UV spectrum of carpachromene (107)	99
3	¹ H NMR spectrum of carpachromene (107)	100
4	¹ H- ¹ H COSY spectrum of carpachromene (107)	101
5	¹³ C NMR and DEPT spectra of carpachromene (107)	102
6	EIMS spectrum of carpachromene (107)	103
7	HMQC spectrum of carpachromene (107)	104
8	IR spectrum of cycloartocarpesin (108)	105
9	¹ H NMR spectrum of cycloartocarpesin (108)	106
10	¹ H- ¹ H COSY spectrum of cycloartocarpesin (108)	107
11	UV spectrum of cycloartocarpesin (108)	108
12	¹³ C NMR and DEPT spectra of cycloartocarpesin (108)	109
13	HMQC spectrum of cycloartocarpesin (108)	110
14	EIMS spectrum of cycloartocarpesin (108)	111
15	HMBC spectrum of cycloartocarpesin (108)	112
16	Infrared spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'',11'',12'':6,7,3',4']flavone (106)	113
17	HREIMS spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'',11'',12'':6,7,3',4']flavone (106)	114
18	HREIMS spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'',11'',12'':6,7,3',4']flavone (106) – continue	115
19	¹ H NMR spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'',11'',12'':6,7,3',4']flavone (106)	116
20	¹ H- ¹ H COSY spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'',11'',12'':6,7,3',4']flavone (106)	117

21	¹³ C NMR spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'',11'',12'':6,7,3',4']flavone (106)	118
22	DEPT spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'', 11'',12'': 6,7,3',4']flavone (106)	119
23	HMQC spectrum of 5-hydroxy-2'',2''-dimethylpyrano- [5'',6'', 11'',12'': 6,7,3',4']flavone (106)	120
24	UV spectrum of 2'-hydroxy-4,4',6'-trimethoxychalcone (109)	121
25	Infrared spectrum of 2'-hydroxy-4,4',6'- trimethoxychalcone (109)	122
26	¹ H NMR spectrum of 2'-hydroxy-4,4',6'- trimethoxychalcone (109)	123
27	¹ H- ¹ H COSY spectrum of 2'-hydroxy-4,4',6'- trimethoxychalcone (109)	124
28	¹³ C NMR and DEPT spectra of 2'-hydroxy-4,4',6'- trimethoxychalcone (109)	125
29	HMQC spectrum of 2'-hydroxy-4,4',6'-trimethoxychalcone (109)	126
30	HMBC spectrum of 2'-hydroxy-4,4',6'-trimethoxychalcone (109)	127
31	EIMS spectrum of 2'-hydroxy-4,4',6'-trimethoxychalcone (109)	128
32	Infrared spectrum of norartocarpetin (48)	129
33	EIMS spectrum of norartocarpetin (48)	130
34	¹ H NMR spectrum of norartocarpetin (48)	131
35	¹ H- ¹ H COSY spectrum of norartocarpetin (48)	132
36	UV spectrum of norartocarpetin (48)	133
37	¹³ C NMR and DEPT spectra of norartocarpetin (48)	134
38	HMQC spectrum of norartocarpetin (48)	135
39	HMBC spectrum of norartocarpetin (48)	136
40	GC-MS spectrum of friedelin (110)	137
41	Infrared spectrum of friedelin (110)	138
42	¹ H NMR spectrum of friedelin (110)	139
43	¹³ C NMR spectrum of friedelin (110)	140
44	¹³ C NMR spectrum of friedelin (110) – expansion	141
45	¹³ C NMR and DEPT spectra of friedelin (110)	142
46	EIMS spectrum of lupeol (111)	143
47	¹ H NMR spectrum of lupeol (111)	144

48	Infrared spectrum of lupeol (111)	145
49	¹³ C NMR and DEPT spectra of lupeol (111)	146
50	EIMS spectrum of β -sitosterol (100)	147
51	Infrared spectrum of β -sitosterol (100)	148
52	¹ H NMR spectrum of β -sitosterol (100)	149
53	¹³ C NMR and DEPT spectra of β -sitosterol (100)	150
54	Infrared spectrum of artonin E (22)	151
55	¹ H NMR spectrum of artonin E (22)	152
56	¹³ C NMR spectrum of artonin E (22)	153
57	¹ H- ¹ H COSY spectrum of artonin E (22)	154
58	DEPT spectrum of artonin E (22)	155
59	HMQC spectrum of artonin E (22)	156
60	UV spectrum of artonin E (22)	157
61	EIMS spectrum of artonin E (22)	158
62	Infrared spectrum of artobiloxanthone (21)	159
63	UV spectrum of artobiloxanthone (21)	160
64	EIMS spectrum of artobiloxanthone (21)	161
65	¹ H NMR spectrum of artobiloxanthone (21)	162
66	¹ H- ¹ H COSY spectrum of artobiloxanthone (21)	163
67	¹³ C NMR and DEPT spectra of artobiloxanthone (21)	164
68	HMQC spectrum of artobiloxanthone (21)	165
69	HMBC spectrum of artobiloxanthone (21)	166
70	Infrared spectrum of catechin (112)	167
71	UV spectrum of catechin (112)	168
72	¹ H NMR spectrum of catechin (112)	169
73	¹ H- ¹ H COSY spectrum of catechin (112)	170
74	¹³ C NMR and DEPT spectra of catechin (112)	171
75	HMQC spectrum of catechin (112)	172
76	HMBC spectrum of catechin (112)	173
77	EIMS spectrum of catechin (112)	174
78	EIMS spectrum of lupeol 3-acetate (101)	175
79	Infrared spectrum of lupeol 3-acetate (101)	176
80	¹ H NMR spectrum of lupeol 3-acetate (101)	177
81	¹³ C NMR and DEPT spectra of lupeol 3-acetate (101)	178

CHAPTER 1

INTRODUCTION

1.1. General Introduction

Medicine and natural products have been closely linked through the use of traditional medicines for thousands of years. Fossil records date human use of plants as medicines at least to the Middle Paleolithic age some 60,000 years ago [1]. Plants have formed the basis of sophisticated traditional medicine systems which have been used in countries such as China and India [2]. These plant-based systems continue to play an essential role in health care, and it were estimated by the World Health Organization that approximately 80% of the world's inhabitants rely mainly on traditional medicines for their primary health care [3].

Traditional medicine is widespread throughout the world and Malaysia is no exception. Traditional medicine plays an important role in the Malaysian society and is expected to play more important role in the future. Malaysia with its tropical rain forest is richly endowed with flora abounds with plants of diverse nature and uses. Moreover, it has a good potential to be developed into various useful natural products. The past ethnomedical surveys suggest that at least about 20% of the estimated total of higher plants of 15,000 species comprise plants which have been reported to possess medicinal and other therapeutic properties [4]. It offers unlimited opportunity for scientists' who are interested in the search of biologically active compounds from plants. There are many reports of these plants which exhibit interesting biological properties including cytotoxic, antimicrobial [5], anti-inflammatory, antioxidant [6], antiplatelet [7], and anti HIV activities [8].

Clinical, pharmacological, and chemical studies of these natural products, which were derived predominantly from plants, were the basis of most early medicines such as aspirin, digitoxin, morphine, quinine, and pilocarpine [9]. An analysis of the origin of the drugs developed between 1981 and 2002 showed that natural products comprised 28% of all new chemical drugs launched into the market. In addition, 24% of these new chemical drugs were synthetic or natural mimic compounds, based on the study of pharmacophores related to natural products. This phenomenon suggests that natural products are important sources for new drugs and also good lead compounds suitable for further modification during drug development [10].

To date, pharmacologists, microbiologists, botanists, and natural-products chemists are combing for phytochemicals and lead molecule that could be developed for the treatment of various diseases.

1.2. Family Moraceae

Moraceae is a family of flowering plants in the order Urticales. Moraceae is often called the mulberry family or fig family. They are usually woody plants with alternate leaves and presence of milky latex. Most species have multiple fruits due to combination of different flowers [11]. This plant family which is distributed in the tropical and subtropical regions of Asia, comprises of 1,400 species divided among 60 genera. Only 9 genera and 137 species could be found in Malaysia, distributed from lowlands to mountain forests [12]. A few of these genera such as *Morus*, *Ficus*, and *Artocarpus* have received the majority of phytochemical investigations and in some cases directed towards the isolation of biologically active constituents.

1.3 Genus *Artocarpus*

Artocarpus is the most commonly encountered genus, representatives of the Moraceae. The genus *Artocarpus* consists of approximately 50 species, which are

native to the region of Southeast Asia. However, the greatest diversity is in Indonesia, Malaysia, and the Philippines [13]. There are 47 species of *Artocarpus* that could be found in Malaysia including the cultivated plants as showed in **Table 1.1**. *Artocarpus* are evergreen or deciduous small plants in which only 20 species are large monoecious trees, with all parts containing white latex. Economically, the genus of *Artocarpus* is appreciable importance as a source of edible fruits, such as *A. heterophyllus* (jackfruit), *A. integer* (cempedak), and *A. communis* (breadfruit). Many species of *Artocarpus* also yield fairly good timber. Some members of this genus have been used medicinally to treat various diseases such as malaria, fever, dysentery and tuberculosis [14].

Table 1.1: *Artocarpus* Species Found in Malaysia [12]

Species	Local Name
<i>A. anisophyllus</i>	Keledang babi
<i>A. bracteata</i>	Ipoh
<i>A. communis</i>	Sukun
<i>A. dadah</i>	Tampang bulu
<i>A. denisonian</i>	Nangka pipit
<i>A. elasticus</i>	Terap nasi
<i>A. fulvicortex</i>	Tampang gajah
<i>A. glaucus</i>	-
<i>A. gomezianus</i>	Tampang hitam
<i>A. heterophyllus</i>	Nangka
<i>A. hispidus</i>	Temponek
<i>A. integer</i>	Cempedak
<i>A. integer var. silvestris</i>	Bangkok
<i>A. kemandu</i>	Pudu
<i>A. lakoocha</i>	Keledang beruk
<i>A. lanceifolius</i>	Keledang
<i>A. lowii</i>	Miku
<i>A. maingayi</i>	Pudu
<i>A. nitidus</i>	Tampang
<i>A. odoratissima</i>	-
<i>A. penduncularis</i>	Miku
<i>A. rigidus</i>	Temponek
<i>A. scortechinni</i>	Terap hitam
<i>A. teysmanii</i>	-

1.3.1 *Artocarpus fulvicortex* Jarrett

Artocarpus fulvicortex Jarrett is one of the Malaysian rare *Artocarpus* species. *A. fulvicortex* Jarrett is locally known as 'keledang tampang gajah'. It is a medium-sized tree with orange brown or reddish brown bark. *A. fulvicortex* can be found in the lowland forests at Perak, Pahang, Negeri Sembilan, Melaka, and Terengganu [12]. However, there is no information on the traditional and medicinal usage of this plant.

1.4 Problem Statement

The *Artocarpus* species is noted as an abundant source of isoprenylated flavonoids, a class of compounds which has the chemical and biological properties attraction. Thus, during the last couple of years, many researches on *Artocarpus* species have been carried out especially by researchers from Japan, Taiwan, and Indonesia. Many phenolic constituents have been discovered and characterized. These metabolites were also significantly bioactive. This information throws a vivid light on *Artocarpus* as medicinal plants. However, there are only few phytochemical and biological activity studies on Malaysian *Artocarpus* species. Within the scope of continuation in search for bioactive compounds from natural plants, this research will focus on the phytochemical and biological activity studies of *Artocarpus fulvicortex* Jarrett.

1.5 Research Objectives

The objectives of this research are to determine the chemical constituents of *A. fulvicortex* which involves isolation and characterization of phytochemicals as well as bioactivity screening since there is no report on the phytochemical study of *A. fulvicortex* Jarrett.

1.6 Scope of Study

This research will focus on two parts of *A. fulvicortex* Jarrett plant which are the leaves and the barks. Several approaches will be endeavoured in order to determine the chemical constituents of *A. fulvicortex* Jarrett. Basically, the dried samples of leaves and barks of *A. fulvicortex* Jarrett will be extracted by using different polarity of organic solvents at room temperature. The crude extracts obtained after removal of solvents will be fractionated into several fractions based on the polarity using vacuum column chromatographic technique. The chemical constituents of each fraction will be analyzed using thin layer chromatography and will then be purified by using gravity column chromatography and recrystallisation to obtain the pure compounds. The structures of the pure compounds will be characterized on the basis of spectroscopic methods including MS, IR, ^1H NMR, ^{13}C NMR, 2D NMR (COSY, HMQC, HMBC) and UV.

The evaluation of the biological activities of the crude extracts and pure compounds will be carried out by using several bioassays. The crude extracts and the pure compounds will be screened for antibacterial, antioxidant, and cytotoxic activities. The antibacterial activity will be tested using disc diffusion method with strains of Gram +ve; *Bacillus subtilis*, *Staphylococcus aureus* and Gram –ve bacteria; *Pseudomonas aeruginosa* and *Escherichia coli*. The antioxidant activity will be performed by using 2,2-diphenyl-1-picrylhydrazyl (DPPH) free radical scavenging assay and measured by UV spectrophotometer. While, 3-(4, 5-dimethylthiazol-2-yl)-2-5-diphenyltetrazolium bromide (MTT) assay was used to evaluate the cytotoxic activity.

REFERENCES

1. Fabricant, D. S. and Farnsworth, N. R. The Value of Plants Used in Traditional Medicine for Drug Discovery. *Environmental Health Perspectives*, 2001. 109: 69-75.
2. Newman, D. J., Cragg, G. M., Snader, K. M. The influence of Natural Products upon Drug Discovery. *Nat. Prod. Rep.*, 2000. 17: 215-234.
3. Mohammad, S. Anticancer Agents from Medicinal Plants. *Bangladesh J. Pharmacol.*, 2006. 1: 35-41.
4. Burkill, I. H. A Dictionary of the Economic Products of the Malay Peninsula. Ministry of Agriculture and Cooperatives: Kuala Lumpur. 1966.
5. Pecchio, M., Solis, P. N., Perez, J. L. L., Vasquez, Y., Rodriguez, N., Olmedo, D., Correa, M., Feliciano, A. S., Gupta, M. P. Cytotoxic and Antimicrobial Benzophenones from the Leaves of *Tovomita longifolia*. *J. Nat. Prod.*, 2006. 69: 410-413.
6. Shin, S. C., Hwang, H. J., Kang, K. J., Lee, B. H. An Antioxidative and Antiinflammatory Agent for Potential Treatment of Osteoarthritis from *Ecklonia cava*. *Arch. Pharm. Res.*, 2006. 29(2): 165-171.
7. Lin, C. N. and Lu C. M. Novel Antiplatelet Constituents from Formosan Moraceous Plants. *J. Nat. Prod.*, 1996. 59: 834-838.
8. Tanaka, N., Okasaka, M., Ishimaru, Y., Takaishi, Y., Sato, M., Okamoto, M., Oshikawa, T., Ahmed, S. U., Consentino, L. M., Lee, K. H. Biyouyanagin A, An Anti-HIV Agent from *Hypericum chinense* L. var. *salicifolium*. *Org. Lett.*, 2005. 7(14): 2997-2999.
9. Butler, M. S. The Role of Natural Product Chemistry in Drug Discovery. *J. Nat. Prod.*, 2004. 67: 2141-2153.

10. Chin, Y. M., Balunas, M. J., Chai, H. B., Kinghorn, A. D. Drug Discovery from Natural Sources. *AAPS J.*, 2006. 8(2): 239-253.
11. Castro, R. M. and Rapini, A. Four New Species of Moraceae from Bahia, Brazil. *Neodiversity*, 2006. 1: 13-20.
12. Whitmore, T. C. Tree Flora of Malaya. Vol 3. Longman Malaysia Sdn. Bhd: Kuala Lumpur. 1978.
13. Hakim, E. H., Juliawaty, L. D., Syah, Y. M., Achmad, S. A. Molecular diversity of *Artocarpus champeden* (Moraceae): A species endemic to Indonesia. *Molecular Diversity*, 2005. 9: 149-158.
14. Hakim, E. H., Asnizar, Yurnawilis, Aimi, N., Kitajima, M., Takayama, H. Artoindonesianin P, A New Prenylated Flavone with Cytotoxic Activity from *Artocarpus lanceifolius*. *Fitoterapia*, 2002. 73: 668-673.
15. Venkataraman, K. Wood Phenolics in the Chemotaxonomy of the Moraceae. *Phytochemistry*, 1972. 11: 1571-1586.
16. Nomura, T., Hano, Y., Aida, M. Isoprenoid-substituted flavonoids from *Artocarpus* plants (Moraceae). *Heterocycles*, 1998. 47: 1179-1205.
17. Nomura, T. The Chemistry and Biosynthesis of Isoprenylated Flavonoids from Moraceous Plants. *Pure Appl. Chem.*, 1999. 71: 1115-1118.
18. Hakim, E. H., Achmad, S. A., Juliawaty, L. D., Makmur, L., Syah, Y. M., Aimi, N., Kitajima, M., Takayama, H., Ghisalberti, E. L. Prenylated Flavonoids and Related Compounds of the Indonesian *Artocarpus* (Moraceae). *J. Nat. Med.*, 2006. 60: 161-184.
19. Kijjoo, A., Cidade, H. M., Pinto, M. M. M., Gonzalez, M. J. T. G., Anantachoke, C., Gedris, T. E., Herz, W. Prenylflavonoids from *Artocarpus elasticus*. *Phytochemistry*, 1996. 43(3): 691-694.
20. Kijjoo, A., Cidade, H. M., Gonzalez, M. J. T. G., Afonso, C. M., Silva, A. M. S., Herz, W. Further Prenylflavonoids from *Artocarpus elasticus*. *Phytochemistry*, 1996. 47(5): 875-878.
21. Mustapha, I., Juliawaty, L. D., Syah, Y. M., Hakim, E. H., Latip, J., Ghisalberti, E. L. An Oxepinoflavone from *Artocarpus elasticus* with

- Cytotoxic activity Against P-338 Cells. *Arch. Pharm. Res.*, 2009. 32(2): 191-194.
22. Ko, H. -H., Lu, Y. -H., Yang, S. -Z., Won, S. -J., Lin, C. -N. Cytotoxic Prenylflavonoids from *Artocarpus elasticus*. *J. Nat. Prod.*, 2005. 68 (11): 1692 -1695.
 23. Syah, Y. M., Achmad, S. A., Ghisalberti, E.L., Hakim, E. H., Makmur, L., Mujahidin, D. Artoindonesianins Q-T, Four Isoprenylated Flavones from *Artocarpus champeden* Spreng. (Moraceae). *Phytochemistry*, 2002. 61: 949-953.
 24. Achmad, S. A., Hakim, E. H., Makmur, L., Mujahidin, D., Juliawaty, L. D., Syah, Y. M. Discovery of Natural Products from Indonesian Tropical Rainforest Plants: Chemodiversity of *Artocarpus* (Moraceae). Kluwer Academic/Plenum Publishers: Indonesia. 2002.
 25. Syah, Y. M., Juliawaty, L. D., Achmad, S. A., Hakim, E. H., Ghisalberti, E. L. Cytotoxic Prenylated Flavones from *Artocarpus champeden*. *J. Nat. Med.*, 2006. 60: 308-312.
 26. Widyawaruyanti, A., Subehan, Kalauni, S. K., Awale, S., Nindatu, M., Zaini, N. C., Syafruddin, D., Asih, P. B. S., Tezuka, Y., Kadota, S. New Prenylated Flavones from *Artocarpus champeden*, and Their Antimalarial Activity In Vitro. *J. Nat. Med.*, 2007. 61(4): 410- 413.
 27. Jayasinghe, L., Balasooriya, B. A. I. S., Padmini, W. C., Hara, N., Fujimoto, Y. Geranyl chalcone derivatives with antifungal and radical scavenging properties from the leaves of *Artocarpus nobilis*. *Phytochemistry*, 2004. 65: 1287-1290.
 28. Jayasinghe, U. L. B., Samarakoon, T. B., Kumarihamy, B. M. M., Hara, N., Fujimoto, Y. Four New Prenylated Flavonoids and Xanthones from the Root Bark of *Artocarpus nobilis*. *Fitoterapia*, 2008. 79: 37-41.
 29. Weng, J. -R., Chan, S. -C., Lu, Y. -H., Lin, H. -C., Ko, H. -H., Lin, C. -N. Antiplatelet Prenylflavonoids from *Artocarpus communis*. *Phytochemistry*, 2006. 67: 824-829.

30. Chan, S. –C., Ko, H. –H., Lin, C. –N. New Prenylflavonoids from *Artocarpus communis*. *J. Nat. Prod*, 2003. 66: 427-430.
31. Lin, K. –W., Liu, C. –H., Tu, H. –Y., Ko, H. –H., Wei, B. –L. Antioxidant Prenylflavonoids from *Artocarpus communis* and *Artocarpus elasticus*. *Food Chem.*, 2009. 115: 558-562.
32. Han, A. –R., Kang, Y. –J., Windono, T., Lee, S. K., Seo, E. –K. Prenylated Flavonoids from the Heartwood of *Artocarpus communis* with Inhibitory Activity on Lipopolysaccharide-Induced Nitric Oxide Production. *J. Nat. Prod.*, 2006. 69 (4): 719 -721.
33. Wei, B. –L., Weng, J. –R., Chiu, P. –H., Hung, C. –F., Wang, J. –P., Lin, C. –N. Antiinflammatory Flavonoids from *Artocarpus heterophyllus* and *Artocarpus communis*. *J. Agric. Food Chem.*, 2005. 53: 3867-3871.
34. Fang, S. –C., Hsu, C. –L., Yen, G. –C. Antiinflammatory Effects of Phenolic Compounds Isolated from the Fruits of *Artocarpus heterophyllus*. *J. Agric. Food Chem.*, 2008. 56: 4463-4468.
35. Arung, E. T., Shimizu, K., Kondo, R. Inhibitory Effect of Artocarpanone from *Artocarpus heterophyllus* on Melanin Biosynthesis. *Biol. Pharm. Bull.*, 2006. 29(6): 1966-1969.
36. Jamil, S., Sirat, H. M., Jantan, I., Aimi, N., Kitajima, M. A New Prenylated Dihydrochalcone from the Leaves of *Artocarpus lowii*. *J. Nat. Med.*, 2008. 62: 321-324.
37. Syah, Y. M., Achmad, S. A., Ghisalberti, E. L., Hakim, E. H., Makmur, L., Mujahidin, D. Artoindonesianins G-I, Three New Isoprenylated Flavones from *Artocarpus lanceifolius*. *Fitoterapia*, 2001. 72: 765-773.
38. Boonphong, S., Baramée, A., Kittakoop, P., Puangsombat, P. Antitubercular and Antiplasmodial Prenylated Flavones from the Roots of *Artocarpus altilis*. *Chiang Mai J. Sci.*, 2007. 34(3): 339-344.
39. Worrell, D. B., Carrington, C. M. S., Huber, D. J. Growth, Maturation, and Ripening of Breadfruit, *Artocarpus altilis* (Park.), Fosb. *Scientia Horticulturae*, 1998. 76: 17-28.

40. Shimizu, K., Kondo, R., Sakai, K., Buabarn, S., Dilokkunanant, U. 5 α -Reductase Inhibitory Component from *Artocarpus altilis*. *J. Wood Sci.*, 2000. 46: 385-389.
41. Chen, C. C., Huang, Y. L., Ou, J. C. Three New Prenylflavones from *Artocarpus altilis*. *J. Nat. Prod.*, 1993. 1594-1597.
42. Namdaung, U., Aroonrerk, N., Suksamrarn, S., Danwisetkanjana, K., Saenboonrueng, J., Arjchomphu, W., Suksamrarn, A. Bioactive Constituents of the Root Bark of *Artocarpus rigidus* subsp. *rigidus*. *Chem. Pharm. Bull.*, 2006. 54(10): 1433-1436.
43. Puntumchai, A., Kittakoop, P., Rajviroongit, S., Vimuttipong, S., Likhitwitayawuid, K., Thebtaranonth, Y. Lakoochins A and B, New Antimycobacterial Stilbene Derivatives from *Artocarpus lakoocha*. *J. Nat. Prod.*, 2004. 67: 485-486.
44. Joshee, N., Bastola, D. R., Agrawal, B. P., Yadav, A. K. Lakoocha: A Multipurpose Tree of Warm Climate. Alexandria: AHS Press. 2002.
45. Hakim, E. H., Ulinuha, U. Z., Syah, Y. M., Ghisalberty, E. L. Artoindonesianins N and O, New Prenylated Stilbene and Prenylated Arylbenzofuran Derivatives from *Artocarpus gomezianus*. *Fitoterapia*, 2002. 73: 597-603.
46. Likhitwitayawuid, K. and Sritularak, B. A New Dimeric Stilbene with Tyrosinase Inhibitory Activity from *Artocarpus gomezianus*. *J. Nat. Prod.*, 2001. 64: 1457-1459.
47. Likhitwitayawuid, K. Stilbenes with Tyrosinase Inhibitory Activity. *Current Sci.*, 2008. 94(1): 44-52.
48. Amarasinghe, N. R., Jayasinghe, L., Hara, N., Fujimoto, Y. Chemical Constituents of the Fruits of *Artocarpus altilis*. *Bio. Syst.s and Ecol.*, 2008. 36: 323-325.
49. Patil, A. D., Freyer, A. J., Killmer, Lew., Offen, P., Taylor, P. B., Votta, B. J., Johnson, R. K. A New Dimeric Dihydrochalcone and A New Prenylated Flavone from the Bud Covers of *Artocarpus altilis*: Potent Inhibitors of Cathepsin K. *J. Nat. Prod.*, 2002. 65:624-627.

50. Shimizu, K., Kondo, R., Sakai, K. A Stilbene Derivative from *Artocarpus incisus*. *Phytochemistry*, 1997. 45(6): 1297-1298.
51. Shimizu, K., Kondo, R., Sakai, K., Sato, H. The inhibitory components from *Artocarpus incisus* on Tyrosinase and Melanin Biosynthesis. 2nd International Electronic Conference on Synthetic Organic Chemistry (ECSOC-2). 1998.
52. Boonlaksiri, C., Oonant, W., Kongsaree, P., Kittakoo, P., Tanticharoen, M., Thebtaranonth, Y. An Antimalarial Stilbene from *Artocarpus integer*. *Phytochemistry*, 2000. 54: 415-417.
53. Jayasinghe, L., Rupasinghe, G. K., Hara, N., Fujimoto, Y. Geranylated Phenolic Constituents from the Fruits of *Artocarpus nobilis*. *Phytochemistry*, 2006. 67: 1353-1358.
54. Fang, S. -C., Hsu, C. -L., Yu, Y. -S., Yen, G. -C. Cytotoxic Effects of New Geranyl Chalcone Derivatives Isolated from the Leaves of *Artocarpus communis* in SW 872 Human Liposarcoma Cells. *J. Agric. Food Chem.*, 2008. 56: 8859-8868.
55. Radwan, M. M., Guzman, R. R., Manly, S. P., Jacob, M., Ross, S. A. Sepicanin A – A New Geranyl Flavanone from *Artocarpus sepicanus* with Activity against Methicillin-Resistant *Staphylococcus aureus* (MRSA). *Phytochemistry*, 2009. (In Press).
56. Wang, Y., Xu, K., Lin, L., Pan, Y., Zheng, X. Geranyl Flavonoids from the Leaves of *Artocarpus altilis*. *Phytochemistry*, 2007. 68: 1300-1306.
57. Shyamamma, S., Chandra, S. B. C., Hedge, M., Naryanswamy, P. Evaluation of Genetic Diversity in Jackfruit (*Artocarpus heterophyllus* Lam.) Based on Amplified Fragment Length Polymorphism Markers. *Genet. Mol. Res.*, 2008. 7(3): 645-656.
58. Chung, M. -I., Lu, C. -M., Huang, P. -L., Lin, C. -N. Prenylflavonoids of *Artocarpus heterophyllus*. *Phytochemistry*, 1995. 40(4): 1279-1282.
59. Jagadeesh, S. L., Reddy, B. S., Swamy, G. S. K., Gorbai, K., Hegde, L., Raghavan, W. G. Chemical Composition of Jackfruit (*Artocarpus*

- heterophyllus* Lamk.) Selections of Western Ghats of India. *Food Chem.*, 2007. 102: 361-365.
60. Shinomiya, K., Aida, M., Hano, Y., Nomura, T. A Diels-Alder Type Adduct from *Artocarpus heterophyllus*. *Phytochemistry*, 1995. 40: 1317-1319.
 61. Langenheim, J. H. Higher Plant Terpenoids: A Phytocentric Overview of Their Ecological Roles. *J. Chem. Ecol.*, 1994. 20(6): 1223-1280.
 62. Mahato, S. B., Banerjee, S. K., Chakravarti, R. N. Triterpenes of the Stem Bark of *Artocarpus chaplasha*. *Phytochemistry*, 1971. 10(6): 1351-1354.
 63. Achmad, S. A., Hakim, E. H., Juliawaty, L. D., Makmur, L., Suyatno. A New Prenylated Flavone from *Artocarpus champeden*. *J. Nat. Prod.*, 1996. 59: 878-879.
 64. Ata, A., Bosch, S. A. V. D., Harwanik, D. J., Pidwinski, G. E. Glutathione S-transferase and Acetylcholinesterase-Inhibiting Natural Products from Medicinally Important Plants. *Pure Appl. Chem.*, 2007. 79(12): 2269-2276.
 65. Heller, W., and Forkmann, G., Biosynthesis. In *The Flavonoids: Advances in Research since 1980*, Harborne, J. B. (ed), Chapman and Hall, London, 399-425, 1988.
 66. Jain, A. C., Khazanchi, R., Kumar, A. Synthesis of Carpachromene. *Chem. Lett.*, 1978. 991-994.
 67. Markham, K. R. Techniques of Flavonoid Identification. London: Academic Press. 1982.
 68. Zheng, Z. -P., Cheng, K. -W., To, J. T. -K., Li, H., Wang, M. Isolation of Tyrosinase Inhibitors from *Artocarpus heterophyllus* and Use of Its Extract as Antibrowning Agent. *Mol. Nutr. Food Res.*, 2008. 52: 1530-1538.
 69. Monache, D. M., Rosa, M. C. D., Scurria, R., Vitali, A., Cuteri, A., Monacelli, B., Pasqua, G., Botta, B. Comparison Between Metabolite Productions in Cell Culture and in Whole Plant of *Maclura pomifera*. *Phytochemistry*, 1995. 39(3): 575-580.
 70. Seidel, V., Bailleul, F., Waterman, P. G. (*Rel*)-1b,2a-di-(2,4-dihydroxy-6-methoxybenzoyl)-3b,4a-di-(4-methoxyphenyl)-cyclobutane and Other

- Flavonoids from the Aerial Parts of *Goniothalamus gardneri* and *Goniothalamus thwaitesii*. *Phytochemistry*, 2000. 55: 439-446.
71. Kishore, P. H., Reddy, M. V. B., Reddy, M. K., Gunasekar, D., Caux, C., Bodo, B. Flavonoids from *Andrographis lineate*. *Phytochemistry*, 2003. 63: 457-461.
 72. Jeong, S. H., Ryu, Y. B., Long, M. J. C., Ryu, H. W., Baek, Y. S., Kang, J. E., Lee, W. S., Park, K. H. Tyrosinase Inhibitory Polyphenols from Roots of *Morus lhou*. *J. Agric. Food Chem.*, 2009. 57: 1195-1203.
 73. Lin, C. -N., Lu, C. -M., Huang, P. -L. Flavonoids from *Artocarpus heterophyllus*. *Phytochemistry*, 1995. 39(6): 1447-1451.
 74. Su, B. -N., Cuendet, M., Hawthorne, M. E., Kardono, L. B. S., Riswan, S., Fong, H. H. S., Mehta, R. G., Pezzuto, J. M., Kinghorn, A. D. Constituents of the Barks and Twigs of *Artocarpus dadah* with Cyclooxygenase Inhibitory Activity. *J. Nat. Prod.*, 2002. 65: 163-169.
 75. Queiroga, C. L., Silva, G. F., Dias, P. C., Possenti, A., Carvalho, J. E. Evaluation on the Antiulcerogenic Activity of Friedelin-3 β -ol and Friedelin Isolated from *Maytenus ilicifolia* (Celastraceae). *J. Ethnopharmacol.*, 2000. 72: 465-468.
 76. Thanakijcharoenpath, W. and Theanphong, O. Triterpenoids from the Stem of *Diospyros glandulosa*. *Thai. J. Pharm. Sci.*, 2007. 31: 1-8.
 77. Subhadhirasakul, S. and Pechpongs, P. A Triterpenoid and Two Steroids from the Flowers of *Mammea siamensis*. *Songklanakarin J. Sci. Technol.*, 2005. 27(2): 556-561.
 78. Agarwal, R. B., and Rangari, V. D. Antiinflammatory and Antiarthritic Activities of Lupeol Isolated from *Strobilanthes callosus* and *strobilanthes ixiocephala* Roots. *Indian J. Pharmacol.*, 2003. 35: 384-387.
 79. John, I. O., and Alexander, E. I. Friedelanone and Other Triterpenoids from *Hymenocardia acida*. *International J. Physic. Sci.*, 2008. 3(6): 156-158.
 80. Fotie, J., Bohle, D. S., Leimanis, M. L., Georges, E., Rukunga, G., Nkengfack, A. E. Lupeol Long-Chain Fatty Acids Esters with Antimalarial Activity from *Holarrhena floribunda*. *J. Nat. Prod.*, 2006. 69: 62-67.

81. Dhakal, R. C., Rajbhandari, M., Kalauni, S. K., Awale, S., Gewali, M. B. Phytochemical Constituents of the Barks of *Vitex negundo* L. *J. Nepal Chem. Soc.*, 2008. 23: 89-92.
82. Jamil, S. Phytochemical and Biological Studies of *Artocarpus lowii* King., *A. scortechinii* King. and *A. teysmanii* Miq. from Malaysia. Faculty Science, UTM (Skudai). PhD Thesis. 2006.
83. Suhartati, T. The Bioactivity Test of Artonin E from the Bark of *Artocarpus rigida* Blume. *European J. Sci. Res.*, 2008. 23(2): 330-337.
84. Suhartati, T., Achmad, S. A., Aimi, N., Hakim, E. H., Kitajima, M., Takayama, H., Takeya, K. Artoindonesianin L, A New Prenylated Flavone with Cytotoxic Activity from *Artocarpus rotunda*. *Fitoterapia*, 2001. 72: 912-918.
85. Alimoza, D. F., Kuliev, Z. A., Vdovin, A. D. Catechins and Proanthocyanidines from *Alhagi pseudoalhagi*. *Chem. Nat. Comp.*, 2007. 43(7): 326.
86. Li, D. -L., Li, X. -M., Peng, Z. -Y., Wang, B. -G. Flavanol Derivatives from *Rhizophora stylosa* and Their DPPH Radical Scavenging Activity. *Molecules*, 2007. 12: 1163-1169.
87. Achmad, S. A., Murniana., Udjiana, S. S., Aimi, N., Hakim, E. H., Makmur, L. Tiga Senyawa Flavan-3-ol dari Tumbuhan *Artocarpus reticulatus*. Institut Teknologi Bandung. 1998.
88. Mishra, P. M., and Sree, A. Chemical Investigation and Antibacterial Study of Hexane Extract of Leaves of *Finlaysonia obovata*. *Cancer Nat. Comp.*, 2009. 45(1): 129-131.
89. Shih, Y. L. and Rothfield, L. The bacterial cytoskeleton. *Microbiol. Mol. Biol. Rev.*, 2006. 70(3): 729-754.
90. Tenover, F. C. Mechanism of Antimicrobial Resistance in Bacteria. *The American J. Med.*, 2006. 119(6): 3-10.
91. Mbaveng, A. T., Ngameni, B., Kuete, V., Simo, I. K., Ambassa, P., Roy, R., Bezabih, M., Etoa, F. X., Ngadjui, B. T., Abegaz, B. M., Meyer, J. J. M., Lall, N. and Beng, V. P. Antimicrobial Activity of the Crude Extracts and

- Five Flavonoids from the Twigs of *Dorstenia barteri* (Moraceae). *J. Ethnopharmacol.*, 2008, 116: 483-489.
92. Arias, M. E., Gomez, J. D., Cudmani, N. M., Vattuone, M. A. and Isla, M. I. Antibacterial Activity of Ethanolic and Aqueous Extracts of *Acacia aroma* Gill. Ex Hook et Arn. *Life Sci.*, 2004. 75: 191-202.
 93. Mori, A., Nishino, C., Enoki, N., Tawata, S. Antibacterial Activity and Mode of Plant flavonoids Against *Proteus vulgaris* and *Staphylococcus aureus*. *Phytochemistry*, 1987. 26:223-224.
 94. Krishnaraju, A. V., Rao, C. V., Rao, T. V. N., Reddy, K. N., Trimurtulu, G. In vitro and In vivo Antioxidant Activity of *Aphanamixis polystachya* Bark. *American J. Infec. Diseases.*, 2009. 5(2): 60-67.
 95. Ko, F. N., Cheng, Z. J., Lin, C. N., Teng, C. M. Scavenger and Antioxidant Properties of Prenylflavones Isolated from *Artocarpus heterophyllus*. *Free Radical Biol. & Med.*, 1998. 25(2): 160-168.
 96. Cory, A. H., Owen, T. C., Barltrop, J. A., Cory, J. G. Use of an Aqueous Soluble Tetrazolium/Formazan Assay for Cell Growth Assays in Culture. *Cancer Comm.*, 1991. 3(7): 207-212.
 97. Mosmann, T. Rapid Colorimetric Assay for Cellular Growth and Survival: Application to Proliferation and Cytotoxicity Assays. *J. Immunological.*, 1983. 65: 55-63.
 98. Tagashira, M. and Ohtake, Y. A. New Antioxidative 1,2-Benzodioxole from *Melissa officianalis*. *Planta Med.*, 1998. 64: 555-558.